



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.


We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let ABC be the given triangle. On AC , AB describe segments to contain angles of 60° . Let H , K be the centers of these circles. Join HK and parallel to HK draw DAE . Join DC and EB , and produce DC , EB until they meet in F . Then FDE is the required maximum equilateral triangle.



$$\angle D = \angle E = \angle F = 60^\circ.$$

Draw any other line GN through A . Let fall the perpendiculars HL , KM , HO , KP , and draw KQ parallel to GN .

$$KQ = OP = \frac{1}{2}GN; HK = LM = \frac{1}{2}DE, HK > KQ. \therefore DE > GN.$$

$\therefore DE$ is the longest line that can be drawn through A .

$\therefore FDE$ is the maximum equilateral triangle required.

The same method will serve to describe the maximum triangle having any angles, about ABC .

Also solved by J. Scheffer.

MISCELLANEOUS.

155. Proposed by A. H. HOLMES, Brunswick, Maine.

There are two vessels, one containing a gallons of alcohol, the other b gallons of water. Suppose that c gallons are simultaneously taken from each and poured into the other, how many times must this be done so that there will be the same proportion of alcohol to water in each vessel?

I. Solution by J. SCHEFFER, A. M., Hagerstown, Md.

Let $F(x)$ represent the number of gallons in the first vessel after x operations, then the number of gallons in the first vessel after the next operation will be $F(x) - \frac{c}{a}F(x) + \frac{c}{b}[a - F(x)]$. $\therefore F(x+1) - [1 - (\frac{c}{a} + \frac{c}{b})]F(x) = \frac{ac}{b}$.

By the calculus of finite differences, we get

$$F(x) = \frac{a^2}{a+b} + C[1 - (\frac{c}{a} + \frac{c}{b})]^x.$$

Since for $x=0$, $F(x)=a$, we get $C = \frac{ab}{a+b}$.

$$\therefore F(x) = \frac{a^2}{a+b} + \frac{ab}{a+b}[1 - (\frac{c}{a} + \frac{c}{b})]^x.$$

It is easily seen that when the same proportion of alcohol to water prevails, the contents of the alcohol in the first vessel will be $=a^2/(a+b)$.

$\therefore x$ must be $=\infty$.

II. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

After the first operation, there are $a-c$ gallons of alcohol in the first vessel, and c gallons of alcohol in the second vessel. After the second operation, there are $a-2c+c^2(1/a+1/b)$ gallons of alcohol in the first vessel, and $2c-c^2(1/a+1/b)$ gallons of alcohol in the second vessel.

Let $A=c(1/a+1/b)$. Then, after the third operation, there are $a-3c+3Ac-A^2c$ gallons of alcohol in the first vessel, and $3c-3Ac+A^2c$ gallons of alcohol in the second vessel. After the n th operation there are

$a-nc+\frac{n(n-1)}{2!}Ac-\frac{n(n-1)(n-2)}{3!}A^2c+\dots\pm A^{n-1}c=a+\frac{c(1-A)^n-c}{A}$ gallons of

alcohol, and $\frac{c-c(1-A)^n}{A}$ gallons of water in the first vessel, and

$nc-\frac{n(n-1)}{2!}Ac+\frac{n(n-1)(n-2)}{3!}A^2c-\dots\pm A^{n-1}c=\frac{c-c(1-A)^n}{A}$ gallons of alco-

hol, and $b+\frac{c(1-A)^n-c}{A}$ gallons of water in the second vessel.

$$\therefore \frac{Aa+c(1-A)^n-c}{c-c(1-A)^n} = \frac{c-c(1-A)^n}{Ab+c(1-A)^n-c}.$$

$$\therefore (1-A)^n = \frac{c(a+b)-Aab}{c(a-b)} = 0. \quad \therefore n = -\infty, \text{ or } A=1.$$

\therefore The result stated can only happen when $a=b=2c$, then $n=1$.

156. Proposed by R. D. CARMICHAEL, Hartselle, Ala.

There exist no multiply perfect odd numbers of multiplicity n containing only n distinct primes.

Solution by JACOB WESTLUND, Ph. D., Purdue University, Lafayette, Ind.

If n denotes the multiplicity of a multiply perfect number $p_1^{a_1} p_2^{a_2} \dots p_i^{a_i}$, where p_1, p_2, \dots are distinct primes, we have

$$n = \frac{p_1 - \frac{1}{p_1^{a_1}}}{p_1 - 1} \cdot \frac{p_2 - \frac{1}{p_2^{a_2}}}{p_2 - 1} \dots, \text{ and hence } n < \frac{p_1}{p_1 - 1} \cdot \frac{p_2}{p_2 - 1} \dots \frac{p_i}{p_i - 1}.$$

Now, if $p_1 > 2$, we have